

Forstliche Forschungsanstalt München

Forschungsberichte

Nr. 9

1973

NEEDLE-BIOMASS, GROWTH AND GROWTH DISTRIBUTION OF PINUS
RADIATA IN SOUTH AFRICA, IN RELATION TO PRUNING AND THINNING

NADEL-BIOMASSE, ZUWACHS UND ZUWACHSVERTEILUNG VON PINUS RADIATA
IN SÜDAFRIKA, UNTER DEM EINFLUSS VON ASTUNG UND DURCHFORSTUNG

FOR-BER

9

von

Anthonie van Laar

Aus dem Institut für Ertragskunde
der Forstlichen Forschungsanstalt München

D-8000 München 40
Amalienstrasse 52

Bibliothek
Institut für
Ertragskunde
München
FV.
Stand: A
Nr. 148

Alle Rechte, insbesondere das Recht der Vervielfältigung und Verbreitung sowie Übersetzung vorbehalten.

Herausgeber: Forstliche Forschungsanstalt München
D-8000 München 40, Amalienstrasse 52

verantwortlich: Der Obmann
der Forstl. Forschungsanstalt
München

Anschrift des Verfassers:

Anthonie van Laar, l.i. (Wag.), D.Sc. (Stell.)

University of Stellenbosch
S t e l l e n b o s c h
South Africa

Zu beziehen über die Universitätsbuchhandlung Heinrich Frank
D-8000 München 40, Amalienstrasse 48

C O N T E N T S

	page
<u>1. INTRODUCTION.</u>	
1.1. The South African silvicultural system in softwood plantations.....	1
1.2. Research objectives.....	7
1.3. Species, soil and climate.....	8
1.4. The water balance.....	12
<u>2. RESEARCH PROCEDURES.</u>	
2.1. Paradyskloof study	17
2.2. Tokai study	23
2.3. Additional studies.....	26
2.4. Statistical procedures.....	27
<u>3. ANALYSIS OF THE PARADYSKLOOF STUDY.</u>	
3.1. Growth per hectare.	
3.1.1. Basal area growth.....	29
3.1.2. Height growth	38
3.1.3. Volume growth.....	41
3.2. Stem analysis.	
3.2.1. Introduction	46
3.2.2. Response during the first year.....	48
3.2.3. Response during the 2nd and 3rd year.....	56
3.2.4. Response during the 4th and 5th year.....	60
3.2.5. Response during the 6th and 7th year.....	66
3.2.6. Discussion.....	70
3.3. Crown morphometric and related studies.	
3.3.1. The relationship between branch diameter and branch length.....	72
3.3.2. Influence of thinning on branch thickness	76
3.3.3. Interrelationships between branch characteristics.....	
3.3.3.1. Relationship between basal area of the primary and basal area of the secondary branches.....	79
3.3.3.2. Relationship between diameter of the primary branch and number of secondary branches.....	81

3.3.3.3. Relationship between diameter of the primary branch and mean diameter of the secondary branches.....	84
3.3.4. Needle dimensions in relation to position within the crown.....	86
3.3.5. The ratio dry weight : fresh weight of the needles in relation to position within the crown.....	94
3.4. Needle-biomass, growth and growth distribution	
3.4.1. The regression model.....	96
3.4.2. Needle weight per tree.	
3.4.2.1. Needle weight of the secondary branch.....	104
3.4.2.2. Needle weight of the primary branch.....	108
3.3.2.3. Total tree needle weight.....	112
3.4.3. Relationship between needle weight and volume growth.....	116
3.4.4. Growth per unit needle weight in relation to position along the stem.....	123
4. ANALYSIS OF THE TOKAI STUDY.	
4.1 Growth per hectare.	
4.1.1. Basal area growth.....	132
4.1.2. Volume growth.....	134
4.2. Relationship between crown size, competition and growth.	
4.2.1. General.....	136
4.2.2. Crown width and crown length in relation to stand density.	
4.2.2.1. Crown width.....	139
4.2.2.2. Crown length.....	140
4.2.2.3. Relationship between crown width and crown length.....	141
4.2.2.4. Relationship between DBH and height above ground of the base of the live crown.....	143
4.2.2.5. The influence of stand density on mean crown width and crown length.....	144
4.2.2.6. Relationship between stem diameter, crown width and tree height.....	147

4.2.2.7. Crown closure.....	150
4.2.3. Growth rate per unit area crown projec-	
tion.	
4.2.3.1. Introduction.....	151
4.2.3.2. Basal area growth.....	152
4.2.3.3. Volume growth.....	155
4.2.4. The influence of intraspecific competi-	
tion within stands.....	156
4.3. Needle-biomass, growth and growth distribution.	
4.3.1. Needle-biomass.	
4.3.1.1. The precision of the estimate..	172
4.3.1.2. Needle-biomass per tree.....	181
4.3.2. Longitudinal growth distribution.....	182
4.3.3. Growth and growth distribution in	
relation to needle-boimass.	
4.3.3.1. Variability of radial growth...	185
4.3.3.2. Basal area growth in relation	
to needle-biomass.....	187
4.3.3.3. Volume growth in relation to	
needle-biomass.....	190
4.4. Crown morphometric studies.	
4.4.1. The distribution of branch parameters....	191
4.4.2. Relation between branch diameter,	
crown section and branch length.....	201
4.4.3. Basal area of the secondary branches	
in relation to diameter, length and	
crown section of the primary branch.....	205
5. MISCELLANEOUS STUDIES, RELATED TO RESPONSE BEHAVIOUR.	
5.1. Early response of <i>P.radiata</i> to pruning.	
5.1.1. Procedure.....	212
5.1.2. Diurnal fluctuations.....	212
5.1.3. Daily growth.....	220
5.2. A crown release experiment.....	225
5.3. The influence of rainfall on growth rate.....	227
5.4. Seasonal periodicity of radial growth.....	233
5.5. Daily radial growth of <i>P.radiata</i> in relation to	
environment.....	240
SUMMARY.....	250
ZUSAMMENFASSUNG.....	258
LITERATURE.....	273
LIST OF TABLES.....	278
LIST OF FIGURES.....	280